

Antimicrobial Stewardship: Tracking, Reporting, Education



Alex Neukirch, PharmD
Consultant Pharmacist
Consonus Pharmacy



Nebraska
Infection Control
Network

1

Key Points

- Outline components recommended by CDC for tracking, reporting, and education
- Provide specific examples of parameters and metrics for tracking
- Review strategies to report antimicrobial stewardship-related activities and outcomes
- Present methods to provide antimicrobial stewardship-related education


2

The Performance Improvement Cycle

Why we track Metrics

What are we trying to Improve?

- Appropriate Antibiotic use
- Resident quality of care




3

Tracking Core Element

4


What Should be Tracked?



<https://www.cdc.gov/longtermcare/pdfs/core-elements-antibiotic-stewardship.pdf>

5

Tracking Antimicrobial Use



6

Why Per 1000 Resident Days?



- ▶ Converts the number of starts, or days of therapy to a rate
- ▶ Allows comparison between facilities of differing size
- ▶ Recommend to do by month

Example:

- ▶ $(\# \text{ of antibiotic starts in August} \div \# \text{ of resident days August}) \times 1000$
- ▶ $(\# \text{ of days of therapy in August} \div \# \text{ of resident days August}) \times 1000$

7

Why Track Standardized Starts and Days of Therapy?



8

What Antimicrobials Should be Tracked?

- ▶ Search “CDC NHSN AUR MODULE”
- ▶ CDC document on how to submit antibiotic use data to NHSN AUR module
- ▶ Provides comprehensive list of antimicrobials
- ▶ Classifies as Anti-viral, Anti-bacterial, or Anti-fungal
- ▶ Provides Drug Class information
- ▶ Only lists generic names

NHSN
NATIONAL HEALTHCARE SAFETY NETWORK
January 2023

Antimicrobial Use and Resistance (AUR) Module

Contents

Antimicrobial Use and Resistance (AUR) Module 1

Introduction 1

1. Antimicrobial Use (AU) Option 2

Introduction 2

Requirements 3

Data Analyses 8

References 16

Appendix A. Table of Instructions: Antimicrobial Use Option 17

Appendix B. List of Antimicrobials 19

Appendix C. Example Calculations of Antimicrobial Days 23

Appendix D. List of SAARs¹ 26

Appendix E. Antimicrobial Groupings for SAAR & Rate Table Calculations² 29

2. Antimicrobial Resistance (AR) Option 36

Introduction 36

Requirements 37

Data Analyses 45

References 54

Appendix F. List of Eligible Organisms for the NHSN AR Option 55

Appendix G. Technical and Isolate Based Report Variables 62

Appendix H. Denominator Data Variables 66

Appendix I. NHSN AR Option Phenotype Definitions 69

Appendix J. List of Skills and Goals 73

Introduction

This module contains two options: one focused on antimicrobial use and the second on antimicrobial resistance. To participate in either option, facility personnel responsible for reporting antimicrobial use (AU) or resistance (AR) data to the National Healthcare Safety Network (NHSN) must coordinate with their pharmacy and/or laboratory information software providers to configure their system to generate standard formatted files to be imported into NHSN. The format provided for data submission follows:

<https://www.cdc.gov/nhsn/pdfs/pscreport/11pscaurcurrent.pdf> Accessed 9/21/23

9

Examples of Antimicrobials for Tracking

Antimicrobial Agent	Antimicrobial Category	Antimicrobial Class ^a	Antimicrobial Subclass ^a	Antimicrobial Agent	Antimicrobial Category	Antimicrobial Class ^a	Antimicrobial Subclass ^a
AMANTADINE	Anti-influenza	M2 ion channel inhibitors		CEFDIOXIME	Antibacterial	Cephalosporins	Cephalosporin 3 rd generation
AMIKACIN	Antibacterial	Aminoglycosides		CEFPROZIL	Antibacterial	Cephalosporins	Cephalosporin 2 nd generation
AMIKACIN LIPOSOMAL ^b	Antibacterial	Aminoglycosides		CEFTAROLINE	Antibacterial	Cephalosporins	Cephalosporins with anti-MRSA activity
AMOXICILLIN	Antibacterial	Penicillins	Aminopenicillin	CEFTAZIDIME	Antibacterial	Cephalosporins	Cephalosporin 3 rd generation
AMOXICILLIN/CLAVULANATE	Antibacterial	β-lactam/β-lactamase inhibitor combination		CEFTAZIDIME/AVIBACTAM	Antibacterial	β-lactam/β-lactamase inhibitor combination	
AMPHOTERICIN B	Antifungal	Polyenes		CEFTOLOZANE/TAZOBACTAM	Antibacterial	β-lactam/β-lactamase inhibitor combination	
AMPHOTERICIN B LIPID COMPLEX	Antifungal	Polyenes		CEFTRIAZONE	Antibacterial	Cephalosporins	Cephalosporin 3 rd generation
AMPHOTERICIN B LIPOSOMAL	Antifungal	Polyenes		CEFURXIME	Antibacterial	Cephalosporins	Cephalosporin 2 nd generation
AMPICILLIN	Antibacterial	Penicillins	Aminopenicillin	CEPHELEXIN	Antibacterial	Cephalosporins	Cephalosporin 1 st generation
AMPICILLIN/SULBACTAM	Antibacterial	β-lactam/β-lactamase inhibitor combination		CHLORAMPHENICOL	Antibacterial	Phenolics	
ANIDULAFUNGIN	Antifungal	Echinocandins		CIPROFLOXACIN	Antibacterial	Fluoroquinolones	
AZITHROMYCIN	Antibacterial	Macrolides		CLARITHROMYCIN	Antibacterial	Macrolides	
AZTREONAM	Antibacterial	Monobactams		CLINDAMYCIN	Antibacterial	Lincosamides	
				COLISTIMETHATE	Antibacterial	Polymyxins	
				COLISTIN [®]	Antibacterial	Polymyxins	
				DALBAVANCIN	Antibacterial	Glycopeptides	Lipoglycopeptides

<https://www.cdc.gov/nhsn/pdfs/pscreport/11pscaurcurrent.pdf> Accessed 9/21/23

10

Drugs That Should NOT Be Counted

- ▶ Antivirals
- ▶ Topical antifungals
 - Nystatin, clotrimazole, ketoconazole
- ▶ Topical antibiotics
 - Triple antibiotic, bacitracin, mupirocin
- ▶ Antibiotic-containing eye and ear drops/ointments
 - Gentamicin, tobramycin, erythromycin
- ▶ Agents that work within GI tract or not absorbed
 - Sulfasalazine, rifaximin
 - Exceptions: vancomycin PO, fidaxomicin
- ▶ Urinary tract antiseptic/analgesic
 - Methenamine, phenazopyridine

11

Determining Antibiotic Starts

- ▶ Obtain data
 - Antibiotic dispense report from pharmacy
 - Quick/Easy to obtain, but may require some fixing (see example)
 - Antibiotic start log
 - More work upfront, but greater control over data

Infection and Antibiotic Start Log Collections

[Infection and Antibiotic Start Log Template](#)



[Infection and Antibiotic Start Log Template Version 2](#)

[Infection and Antibiotic Start Log Template Version 2 \(demo only\)](#)

<https://asap.nebraskamed.com/facilities/long-term-care/tools-and-templates-for-long-term-care/>

12

Determining Antibiotic Starts

Pharmacy Dispense Report Example

May Antibiotic Report

Product Name	Label Name	Days Supply	Quantity Dispensed	Date of Service	The Clinician
SULFAMETHOXAZOLE-TRIMETHOPRIM	SULFAMETHOXAZOLE-TRIM-DS TABLET	5	15	10/17/2016	SULFONAMIDES (SYSTEMIC)
CEPHALEXIN	CEPHALEXIN 500 MG CAPSULE	10	29	1/19/2016	CEPHALOSPORINS
CEPHALEXIN	CEPHALEXIN 500 MG CAPSULE	10	29	1/19/2016	CEPHALOSPORINS
CEPHALEXIN	CEPHALEXIN 250 MG CAPSULE	1	2	1/19/2016	CEPHALOSPORINS
CEPHALEXIN	CEPHALEXIN 250 MG CAPSULE	1	2	1/19/2016	CEPHALOSPORINS
CEFUROXIME	CEFUROXIME AXETIL 500 MG TAB	7	7	12/22/16	CEPHALOSPORINS
CEFUROXIME	CEFUROXIME AXETIL 250 MG TAB	4	4	12/22/16	CEPHALOSPORINS
CEFUROXIME	CEFUROXIME AXETIL 250 MG TAB	1	1	12/16/2016	CEPHALOSPORINS
CEFDIOXIME PROXETIL	CEFDIOXIME 200 MG TABLET	7	14	12/27/2016	CEPHALOSPORINS
CEFDIOXIME	CEFDIOXIME 200MG CAPSULE	1	1	10/10/2016	CEPHALOSPORINS

May Resident-Days = 450

- ▶ Remove agents that shouldn't be counted
 - Bacitracin, clotrimazole
- ▶ Assume each line represent one antibiotic start
 - 9 lines = 9 starts
- ▶ Calculate starts/1000 resident-days

$$\frac{9 \text{ starts}}{450 \text{ resident-days}} * 1000 = 20$$

13

Other Things to Consider

May Antibiotic Dispense Report

Product Name	Label Name	Days Supply	Quantity Dispensed	Date of Service	The Clinician
SULFAMETHOXAZOLE-TRIMETHOPRIM	SULFAMETHOXAZOLE-TRIM-DS TABLET	5	15	10/17/2016	SULFONAMIDES (SYSTEMIC)
CEPHALEXIN	CEPHALEXIN 500 MG CAPSULE	10	29	1/19/2016	CEPHALOSPORINS
CEPHALEXIN	CEPHALEXIN 500 MG CAPSULE	10	29	1/19/2016	CEPHALOSPORINS
CEPHALEXIN	CEPHALEXIN 250 MG CAPSULE	1	2	1/19/2016	CEPHALOSPORINS
CEPHALEXIN	CEPHALEXIN 250 MG CAPSULE	1	2	1/19/2016	CEPHALOSPORINS
CEFUROXIME	CEFUROXIME AXETIL 500 MG TAB	7	7	12/22/16	CEPHALOSPORINS
CEFUROXIME	CEFUROXIME AXETIL 250 MG TAB	4	4	12/22/16	CEPHALOSPORINS
CEFUROXIME	CEFUROXIME AXETIL 250 MG TAB	1	1	12/16/2016	CEPHALOSPORINS
CEFDIOXIME PROXETIL	CEFDIOXIME 200 MG TABLET	7	14	12/27/2016	CEPHALOSPORINS
CEFDIOXIME	CEFDIOXIME 200MG CAPSULE	1	1	10/10/2016	CEPHALOSPORINS

May Resident-Days = 450

- ▶ Duplicate listing with same drug, dose, and duration
 - For different residents?
 - Need patient level data to confirm
 - Charge to different insurance?
 - Need patient level data and prescription number
- ▶ Was any antibiotic ordered, dispensed, but not given?
 - Request pharmacy for drug crediting report if available
 - Review antibiotic start log

14

Determining Antibiotic Days of Therapy

August Antibiotic Report

Product Name	Label Name	Days Supply	Quantity Dispensed	Date of Service	The Class Name
AMOXICILLIN-CLAVULANATE POTASS	AMOXICLAV 400-57 MG/62.5 ML SUSP	7	100	8/1/2016	PENICILLINS
CEFUROXIME	CEFUROXIME AXETIL 250 MG TAB	1	2	8/2/2016	CEPHALOSPORINS
DOXYCYCLINE HYCLATE	DOXYCYCLINE HYCLATE 100 MG TAB	16	16	8/3/2016	TETRACYCLINES
CEPHELEXIN	CEPHELEXIN 500 MG CAPSULE	7	21	8/3/2016	CEPHALOSPORINS
CEFUROXIME	CEFUROXIME AXETIL 250 MG TAB	9	18	8/3/2016	CEPHALOSPORINS
DOXYCYCLINE HYCLATE	DOXYCYCLINE HYCLATE 100 MG TAB	7	14	8/3/2016	TETRACYCLINES
NITROFURANTON	NITROFURANTON MCR 100 MG CAP	1	1	8/3/2016	URINARY ANTI-INFECTIVES
NITROFURANTON MONO-MACRO	NITROFURANTON MONO-MCR 100 MG	7	14	8/3/2016	URINARY ANTI-INFECTIVES
CEPHALOSPORIN	CEPHALOSPORIN 500 MG	6	12	8/3/2016	CEPHALOSPORINS

August Resident-Days = 490

- Obtain data
 - Antibiotic dispensing report
 - Antibiotic start log if duration recorded
- Remove drug that should not be counted (*none*)
- Sum up 'Days Supply' for all antibiotics or for specific classes
 - All agents: 49 days
 - Cephalosporins: 17 days
- Calculate days of therapy (DOT) per 1000 resident-days

$$\text{All} = \frac{49 \text{ days}}{490 \text{ resident-days}} * 1000 = 100$$

Cephalosporins = ??

15

Other Things to Consider

August Antibiotic Report


Product Name	Label Name	Days Supply	Quantity Dispensed	Date of Service	The Class Name
AMOXICILLIN-CLAVULANATE POTASS	AMOXICLAV 400-57 MG/62.5 ML SUSP	7	100	8/1/2016	PENICILLINS
CEFUROXIME	CEFUROXIME AXETIL 250 MG TAB	1	2	8/2/2016	CEPHALOSPORINS
DOXYCYCLINE HYCLATE	DOXYCYCLINE HYCLATE 100 MG TAB	10	20	8/3/2016	TETRACYCLINES
CEPHELEXIN	CEPHELEXIN 500 MG CAPSULE	7	21	8/3/2016	CEPHALOSPORINS
CEFUROXIME	CEFUROXIME AXETIL 250 MG TAB	9	18	8/3/2016	CEPHALOSPORINS
DOXYCYCLINE HYCLATE	DOXYCYCLINE HYCLATE 100 MG TAB	7	14	8/3/2016	TETRACYCLINES
NITROFURANTON	NITROFURANTON MCR 100 MG CAP	1	1	8/3/2016	URINARY ANTI-INFECTIVES
NITROFURANTON MONO-MACRO	NITROFURANTON MONO-MCR 100 MG	7	14	8/3/2016	URINARY ANTI-INFECTIVES
CEPHALOSPORIN	CEPHALOSPORIN 500 MG	6	12	8/3/2016	CEPHALOSPORINS

August Resident-Days = 490

- Was antibiotic prescribed and dispensed but not given at all?
- Was antibiotic stopped earlier than initially prescribed?
- What happens if patient only received 1 dose out of many prescribed for a single day (e.g., doxycycline)?

16


Electronic Infection and Antibiotic Start Log using Microsoft Excel



Available at: <https://asap.nebraskamed.com/long-term-care/tools-templates-long-term-care/>. Accessed 9/21/23.

17

Electronic Infection and Antimicrobial Start Log using Microsoft Excel



Available at: <https://asap.nebraskamed.com/long-term-care/tools-templates-long-term-care/>. Accessed 4/15/2019.

18

Self-Assessment Question #1

A resident was taking azithromycin and amoxicillin for 5 days for community-acquired pneumonia

1. What is the total number of antibiotic starts?
2. What is the total days of therapy?



19

Self-Assessment Question #2

A resident started Nitrofurantoin twice a day for 5 days starting the evening of September 30th, and finished the course on the morning of October 5th.

1. Does the start count for September, or October?
2. What are the total days of therapy for September?
3. What are the total days of therapy for October?



20

Why Track So Many Use Metrics?

Antibiotic Starts

- Assess impact of initiatives that address when antibiotics are not appropriate (e.g., asymptomatic bacteriuria)

Days of Therapy

- Assess impact of interventions that shorten duration of therapy
- Better metric to monitor overall antibiotic use over time

Standardizing by Patient-Days

- Account for variations in number of residents and lengths of stay
- Allow within facility comparison over time
- Make between facility comparison possible


21

Tracking Process Measures – *Compliance with Clinical Assessment*

<u>What to Track</u>	<u>How to Organize Data</u>	<u>How to Present Data</u>
Was assessment tool used? (Y/N)	Overall (200 assessment performed)	% of time assessment tool used for an infection
Which tool was used? (UTI SBAR, RTI SBAR)	By type of infection (UTI, SSTI)	% of suspected infection met criteria before starting ABX
Were criteria met? (Y/N)	By unit (2 east, 2 west)	
Who assessed the resident? (Nurse A, Nurse B)	By person assessing resident (Nurse A, Nurse B)	

22

Examples of Clinical Assessment Tools



23

Tracking Process Measure – Prescribing Documentation

What to Track	<ul style="list-style-type: none"> • Antibiotic orders with dose, frequency, duration, indication documented
How to Organize Data	<ul style="list-style-type: none"> • Overall • By prescribers
How to Present Data	<ul style="list-style-type: none"> • % of all antibiotic orders with required documentation • % of antibiotic orders from a specific prescriber with required documentation
Logistics	<ul style="list-style-type: none"> • Capture data <u>before</u> getting missing info • Request nurses to capture info for all new antibiotic orders

24

Tracking Process Measure – *Antibiotic Selection Based on Guidelines*

- ▶ Only if facility-specific treatment guidelines are available
 - Based on national guidelines, resistance pattern, prescriber preferences
 - Work with consultant pharmacists, medical directors to create
- ▶ What to track
 - Frequency guideline-recommended antibiotics are selected
 - Frequency the correct dose is selected
 - Frequency correct duration is prescribed
- ▶ Capture data monthly or quarterly if antibiotic use is low
- ▶ Data can be from
 - Consultant Pharmacist
 - Antibiotic start/infection log
 - Indication **MUST** be documented for successful tracking



25

Tracking Outcome Measures – *Antibiogram*

- ▶ Antibiotic susceptibility patterns for specific organisms in a period
- ▶ Based on >30 isolates of an organism to increase statistical power
 - But >20 isolates acceptable per AHRQ
 - Can increase isolates by increasing timespan (e.g., to 24 months)
- ▶ Should only base information on 1st positive culture from multiple consecutive positives
- ▶ If large number of positive cultures, can categorize antibiogram
 - By culture source (e.g., urine cultures)
 - By nursing units



26

Other Outcome Measures – Adverse Drug Events (ADR)

- ▶ Rates of antibiotic-related adverse events
 - *C difficile* infections
 - Diarrhea, loose stools unrelated to CDI
 - Rash, hives
 - Fluoroquinolones: Tendon rupture, hypo/hyperglycemia, confusion, seizure, neuropathy, others
- ▶ Requires careful review of clinical records to determine causality
 - Naranjo adverse drug reaction probability scale¹
 - Classify causal relationship as definite, probable, possible, doubtful
- ▶ Request assistance from
 - Consultant pharmacist as part of monthly drug use evaluation
 - Medical director
 - Specific nurse caring for resident who experienced the ADR

Naranjo CA, et al. Clin Pharmacol Ther 1981;30:239-45.

29

Example of Tool for Evaluating ADR

[Facility Logo] Resident Label

Adverse Drug Reaction Worksheet

Evaluation Date: _____ Evaluated by: _____ Date of Adverse Reaction: _____

Subject Medication: _____ Ongoing regimen: _____ Is medication new? Yes No

Adverse Reaction: _____
(Refer to a list of common adverse antimicrobial reactions in Table 1 on the next page)

Probability Reaction Related to Medication (from the Naranjo Probability Scale below):
 Question (1) Possible (2-4) Doubtful (5)

Consequence of Adverse Reaction (check all that apply):
 Therapy changed to another agent Therapy discontinued
 Increased monitoring Symptomatic medical treatment Corrective surgical procedure
 Hospitalization Permanent damage Delayed discharge
 Other (specify) _____

Reason: _____ Date: _____

Naranjo Adverse Drug Reaction Probability Scale (with modifications)

The following scale is used to assess the likelihood a particular adverse reaction is related to a medication. Answer each of the 10 questions, calculate total score, and determine if an adverse drug reaction is definite, probable, possible, or unlikely related to the drug in question. Interpretation of the probability classification can be found in Table 2 on the next page!

Question	Yes	No	No Yes/No	Score
1. Are there previous CONCOMITANT events on this resident?	+1	0	0	
2. Did the adverse reaction appear after the suspected drug was administered?	+2	-1	0	
3. Did the adverse reaction improve when the drug was discontinued or a specific antagonist given?	+1	0	0	
4. Did the adverse reaction reappear when the drug was readministered?	+2	-1	0	
5. Are there alternative causes (other than the suspect drug) that could have caused the reaction?	-1	+2	0	
6. Did the reaction reappear when a placebo was given?	-1	+1	0	
7. Was the drug absent or blood or other fluids in concentrations known to be toxic?	+1	0	0	
8. Was the reaction more severe when the drug was given at a higher than usual dose or over a longer period of time?	+1	0	0	
9. Did the patient have a similar reaction to the same or similar drug in any previous exposure?	+1	0	0	
10. Was the adverse reaction confirmed by any objective evidence?	+1	0	0	
Total Score				

From: <https://asap.nebraskamed.com/long-term-care/tools-templates-long-term-care/>. Accessed 4/15/19.

30

Table 1. List of Common Adverse Antimicrobial Reactions*

Drug Class	Drug Member	Common Adverse Reaction
Penicillins +/- Beta-Lactamase inhibitors	Ampicillin, Ampicillin Sulbactam, Amoxicillin, Amoxicillin-Clavulanate, Cloxacillin, Dicloxacillin, Nafcillin, Oxacillin, Penicillin, Piperacillin-Tazobactam	Nausea, vomiting, diarrhea, <i>C difficile</i> infection, allergic reactions (including rash, hemolytic anemia), elevated serum creatinine, bone marrow suppression with long-term use, phlebitis with IV therapy
	Cephalosporins +/- Beta-Lactamase inhibitors	Cefaclor, Cefazolin, Cefdinir, Cefepime, Cefixime, Cefotaxime, Cefotetan, Cefprozim, Cefuroxime, Ceftriaxone, Cefuroxime-Tazobactam, Cefuroxime, Cefuroxime, Cephalosporins, Cephalosporins
Carbapenems	Ertapenem, Ertapenem, Imipenem-Cilastatin, Meropenem	Nausea, vomiting, diarrhea, <i>C difficile</i> infection, seizure
Fluoroquinolones	Ciprofloxacin, Delamanid, Levofloxacin, Moxifloxacin	Disorientation, delirium, agitation, seizure, hypo- or hyperglycemia, peripheral neuropathy, tendon rupture, QT prolongation, nausea, vomiting, <i>C difficile</i> infection, increased in liver function tests, aortic dissection
Macrolides	Azithromycin, Clarithromycin, Erythromycin	Nausea, vomiting, elevation in liver function tests, reversible tasteless or tasteless, taste alteration, phlebitis with IV therapy
Tetracyclines	Demeclocycline, Minocycline, Tetracycline	Nausea, vomiting, sunburn, esophageal ulcer, phlebitis with IV therapy, teeth discoloration
Sulfonamides	Sulfamethoxazole-Trimethoprim	Allergic reactions (rash, hives, drug fever), Steven Johnson Syndrome, headache, sunburn, hyperkalemia, worsened renal functions, bone marrow suppression, hemolytic anemia, hypoglycemia (especially with sulfonamides)
Glycopeptides	Telavancin, Vancomycin IV	Redman syndrome (flushing, itching, hypotension), worsened renal functions
Others	Clindamycin, Metronidazole, Nitrofurantoin	All: Nausea, vomiting, Clindamycin: Diarrhea, <i>C difficile</i> infection, taste alteration; Metronidazole: disulfiram reaction after alcohol (flushing, dyspepsia), taste alteration, peripheral neuropathy, confusion; Nitrofurantoin: interstitial pneumonitis (especially with chronic use), hemolytic anemia

* The above list does not include all antimicrobials or all adverse drug reactions. Consult drug references and published literature for additional information if an adverse drug reaction not listed above is suspected.

Table 2. Interpretation of Probability Categories

Category	Score Range	Interpretation
Definite	≥ 9	Reaction 1) followed a reasonable temporal sequence after a drug or in which a toxic drug level had been established in body fluids or tissues; 2) followed a recognized response to the suspected drug; and 3) was confirmed by withdrawal but not by exposure to the drug
Probably	5-8	Reaction 1) followed a reasonable temporal sequence after a drug; 2) followed a recognized response to the suspected drug; 3) was confirmed by withdrawal but not by exposure to the drug; 4) could not be reasonably explained by the known characteristics of the patient's clinical staff
Possible	1-4	Reaction 1) followed a temporal sequence after a drug; 2) possibly followed a recognized pattern to the suspected drug; 3) could be explained by characteristics of the patient's disease
Doubtful	0	Reaction was likely related to factors other than a drug

Reference: Naranjo CA, et al. A method for estimating the probability of adverse drug reactions. Clin Pharmacol Ther 1981;30:239-45.

Other Antibiotic Use/Outcome Measures NOT Covered

- ▶ Defined daily dose (DDD) per 1000 resident-days
 - Requires more calculations
 - DDD definitions may not reflect how antibiotics are usually dosed
- ▶ Point prevalence survey
 - % residents receiving antibiotics on a single day
 - Easier to determine but does not inform overall use
- ▶ Antibiotic utilization ratio
 - Total monthly DOT/total monthly resident-days
 - Represent average DOT in a single resident-day
- ▶ Antibiotic-related costs
 - Important from an administrative perspective
 - Costs fluctuate and may not represent overall use

31

Reporting Core Element

32

Reporting Antimicrobial Stewardship Data Activities and Outcomes

- ▶ CDC recommends reporting tracked data to:
 - Clinical providers
 - Nursing staff
- ▶ ASAP experience from onsite visits
- ▶ Tracked data typically only available to a select few (e.g., QAPI)
- ▶ What good is it if no one knows about it!
 - Data can increase buy-in
 - Resistance rate may deter use
 - Rate of inappropriate UA/culture may improve use of assessment tool
 - Justify your existence in the facility

33

Reporting Antimicrobial Stewardship Activities and Outcomes

- ▶ Who to report to
 - Prescribers
 - Nursing staff
 - Selectively to residents/families
- ▶ What to report
 - Antibiotic use data (starts, DOT)
 - Rates of specific infections
 - Compliance to policy (met criteria before starting antibiotics)
 - \$\$ spent on antibiotics
- ▶ How to report
 - Use existing system
 - Newsletter, QAPI report
 - Frequency varies based on type and volume of information
 - Annually for antibiogram, antibiotic spending
 - Quarterly, semi-annually, or annually for other info

34

Examples of Annual ASP Report

Sample Annual ASP Summary Report--For LTCF Staff [Facility Logo]

Sample Annual Antimicrobial Stewardship Program Report

Highlights of Antimicrobial Stewardship Program (ASP) Initiatives in 20xx

- Implemented use of SBAR tool for suspected UTI (completed)
- Revised contract with laboratory to include production of facility-specific antibiogram (completed)
- Revised contract with consultant pharmacy to review all antimicrobial use (completed)
- Provided mandatory antimicrobial stewardship education to all facility staff (completed)
- Successfully surveyed by CMS on IPCP and ASP without receiving an F-tag (completed)
- Implementing use of SBAR tool for suspected respiratory tract infections (ongoing)
- Collaborating with Nebraska ASAP to further augment ASP (ongoing)

Antimicrobial Stewardship Program-Related Outcomes in 20xx

A. Process Measures

- Compliance with antimicrobial prescribing documentation
 - 250 prescriptions written
 - 80% with all the required information (dose, duration, indication)
- Compliance with use of SBAR for suspected UTI
 - 200 SBAR used and 160 (80%) were fully completed

From: <https://asap.nebraskamed.com/long-term-care/tools-templates-long-term-care/>. Accessed 4/15/19.

35

Sample Annual ASP Summary Report--Personalized for LTCF Prescriber, Dr. Prescriber [Facility Logo]

Sample Annual Antimicrobial Stewardship Program Report

Highlights of Antimicrobial Stewardship Program (ASP) Initiatives in 20xx

- Implemented use of SBAR tool for suspected UTI (completed)
- Revised contract with laboratory to include production of facility-specific antibiogram (completed)
- Revised contract with consultant pharmacy to review all antimicrobial use (completed)
- Provided mandatory antimicrobial stewardship education to all facility staff (completed)
- Successfully surveyed by CMS on IPCP and ASP without receiving an F-tag (completed)
- Implementing use of SBAR tool for suspected respiratory tract infections (ongoing)
- Collaborating with Nebraska ASAP to further augment ASP (ongoing)

Antimicrobial Stewardship Program-Related Outcomes in 20xx

A. Process Measures: Compliance to Antimicrobial Prescribing Documentation

- Total Number of Antimicrobial Prescribed: 250
 - Missing 11 required elements: 50 (20%)
 - Over: 5 (2%)
 - Duration: 20 (8%)
 - Indication: 40 (16%)

B. Antimicrobial Use Measures

- Antimicrobial starts/1000 resident-day is 1% lower compared to 20xx
- Antimicrobial days of therapy/1000 resident-day is 7% lower compared to 20xx
- Antimicrobial appropriateness decreased by 10% compared to 20xx

	Antimicrobial Starts / 1000 Resident Day	Days of Therapy / 1000 Resident Day	Appropriateness
20xx	14.17	24.88	121.68
20xx	1.96	1.98	82.54
20xx	1.96	1.57	22.86
20xx	1.38	1.40	13.82

Most Frequently prescribed antimicrobials

Antimicrobial	Starts / 1000 Resident Day	Days of Therapy / 1000 Resident Day	Appropriateness
Penicillins	1.96	1.98	82.54
Fluoroquinolones	1.96	1.57	22.86
Cephalosporins	1.38	1.40	13.82

Comparison of Antimicrobial Starts by Prescribers

Education Core Element

36

Antimicrobial Stewardship Education

Who to Educate	What to Educate	When to Educate	How to Educate
<ul style="list-style-type: none"> • Prescribers • Nurses • Residents • Families 	<ul style="list-style-type: none"> • Appropriate use • Mcgeer/Loeb Criteria • Consequences of misuse • What is ASP • Program goals • Interventions • Program leaders 	<ul style="list-style-type: none"> • As needed • <i>Annually</i>: antibiogram, general ASP education • <i>Quarterly</i>: Resistance / <i>C diff</i> rates, DOT, antibiotic starts • <i>As needed</i>: New residents, employees, interventions 	<ul style="list-style-type: none"> • Pamphlets / Flyers • Pocket guides • Newsletters • Face-to-face interactive lectures • Web-based education

37

The screenshot shows an email template for an Antimicrobial Stewardship Program (ASP) education. It includes a header with a facility logo, a 'FROM' field (Executive Director, Medical Director, etc.), a 'DATE' field, and a 'RE' field (Antimicrobial Stewardship Program). The main body of the email contains several paragraphs of text. Red callout boxes highlight specific sections: 'Consequence of misuse' points to a paragraph about antibiotic misuse leading to undesirable outcomes; 'What is ASP & Goal of ASP' points to a paragraph describing the program's purpose; 'Program leaders' points to a paragraph stating that infection preventionists play a central role; 'Specific ASP tasks' points to a paragraph listing tasks like reviewing antibiotic courses and making recommendations; and 'Who is responsible' points to a paragraph stating that the success of the initiative requires full participation and support from prescribers, staff, residents, and families.

Example of ASP Education

<https://asap.nebraskamed.com/facilities/long-term-care/educational-materials-for-long-term-care/>

38

Prescriber Education

Effective Communication about Respiratory Infections

SCENARIO 2
Mr. Jones woke up with a cough. He is concerned and asks for an antibiotic because in the past, antibiotics have helped him feel better when he is sick.

Antibiotics should not be prescribed for residents with upper respiratory infections or acute uncomplicated bronchitis unless pneumonia is suspected, or they meet criteria for antibiotic initiation?

Healthcare professionals can use the 4-part Communication Strategy* discussed above to avoid unnecessary antibiotic treatment for residents with respiratory tract infections.

- 1. Review findings:**
Mr. Jones, I am sorry you are not feeling well today. When I examined you, your oxygen level and temperature were normal, you have no throat swelling or nasal tenderness, and your lungs sounded clear.
- 2. Deliver a clear diagnosis:**
The doctor and I discussed your symptoms. It seems that you have acute bronchitis, also known as a chest cold, which is most commonly caused by a virus.
- 3. Provide a FIRST negative, THEN positive treatment recommendation:**
An antibiotic will not work against a viral infection, and may cause side effects, such as diarrhea. Instead, we will treat you for respiratory viruses, including flu. We will provide treatment to help you feel better and closely monitor your symptoms.
- 4. Discuss a contingency plan:**
If you become short of breath, develop a fever or any other concerning symptoms, we will perform more testing, a chest x-ray, and start antibiotics if needed.

From: <https://www.cdc.gov/antibiotic-use/pdf/antibiotic-use-toolkit/508.pdf>. Accessed 9/1/24

39

Resident/Family-Specific ASP Educations

What to Ask Your Healthcare Provider about Antibiotics

Core Elements for Antibiotic Stewardship in Nursing Homes

What You Need to Know About Antibiotics in a Nursing Home

What are antibiotics?
Antibiotics are drugs used to treat infections caused by bacteria. They do not work for illnesses caused by viruses, like flu and most cases of bronchitis.

What is antibiotic stewardship?
Antibiotic stewardship refers to a set of commitments and actions designed to make sure patients receive the right drug, at the right antibiotic, for the right amount of time; and only when truly necessary. Improving antibiotic use will ensure these life-saving medications are effective and available when we need them.

When are antibiotics necessary?
There are times when antibiotics are needed.

Are antibiotics needed - stop, proceed with caution, or go?

These information is provided to you by Nebraska Antimicrobial Stewardship Assessment and Promotion Program (ANSAP) and Nebraska Infection Control Assessment and Promotion Program (NICAP).

Decisions about antibiotic treatment should be made during conversations with your antibiotic user. Healthcare providers prescribe antibiotics... There may be some cases to treat your symptoms.

Antibiotics...

When should you take these infection fighting drugs?

Learn to protect yourself and your community

From: <https://www.cdc.gov/antibiotic-use/education/antibiotic-stewardship.html>. Accessed 9/18/17.

40

Resident/Family-Specific ASP Educations – Trifold Brochure

Why does taking antibiotics lead to antibiotic resistance?

Any time you take antibiotics, they can cause side effects and contribute to the development of antibiotic resistance. Antibiotic resistance is one of the most urgent threats to the public's health.

Always remember:

1. Antibiotic resistance does not mean the body is becoming resistant to antibiotics; it means bacteria are developing the ability to defeat the antibiotics designed to kill them.
2. When bacteria become resistant, antibiotics cannot fight them, and the bacteria multiply.
3. Some resistant bacteria can be harder to treat and can spread to other residents in the nursing home.

Up to 70% of residents in a nursing home receive one or more courses of antibiotics each year.



What if I have questions about antibiotics?

Talk to your healthcare professional if you have any questions about your antibiotics, such as:

- What infection does this antibiotic treat and do you know I have that infection?
- How long do I need to take the antibiotic?
- What are the potential side effects from this antibiotic?
- Could any of my other medications interact with this antibiotic?
- How will you know that the antibiotic is working for my infection?

Improving the way healthcare professionals prescribe antibiotics, and the way we take antibiotics, helps keep us healthy now, helps fight antibiotic resistance, and ensures that these life-saving drugs will be available for future generations.

40%-75% of antibiotics prescribed in nursing homes may be unnecessary or inappropriate.

To learn more about antibiotic prescribing and use, visit www.cdc.gov/antibiotic-use or call 1-800-CDC-INFO.



Do You Need Antibiotics?



Why is it important to Be Antibiotics Aware in nursing homes?

Antibiotics are life-saving drugs and are frequently prescribed in nursing homes. Remember, when antibiotics are needed, their benefits outweigh the risks of side effects and antibiotic resistance.

When antibiotics aren't needed, they won't help you, and the side effects could still cause harm.

What do antibiotics treat?

Antibiotics are only needed for treating certain infections caused by bacteria. Antibiotics are critical tools for treating life-threatening conditions such as pneumonia and sepsis, which is the body's extreme response to an infection.

What don't antibiotics treat?

Antibiotics do not work on viruses, such as those that cause colds, flu, bronchitis, or runny noses, even if the mucus is thick, yellow, or green. Antibiotics also won't help some common bacterial infections, including most cases of bronchitis, many sinus infections, and some eye infections.

What are the potential side effects of antibiotics?

- Common side effects from antibiotics can include:
- Rash
 - Nausea
 - Yeast infections
 - Diarrhea
 - Dizziness

More serious side effects can include:

- Life-threatening allergic reactions
- Interactions between antibiotics and other medications
- Infections with antibiotic-resistant bacteria, including C. difficile (or C. diff)

What is a C. diff infection?

C. diff is a bacterial infection that needs immediate treatment. It can cause severe diarrhea that can lead to severe colon damage and death.

Antibiotics fight bacterial infections by killing bad germs, but can also get rid of the good germs that protect your body against harmful infections. The effect of antibiotics can last as long as several months. If you come in contact with C. diff germs during this time, you can get sick.

- Symptoms of a C. diff infection include:
- Severe diarrhea
 - Stomach tenderness or pain
 - Fever
 - Loss of appetite
 - Nausea

More than 3 million Americans receive care or reside in nursing homes every year.

How can I stay healthy?

- You can stay healthy and keep others healthy by:
- Insisting healthcare professionals and visitors clean their hands before touching you by washing with soap and water for 20 seconds or using a hand sanitizer that contains at least 60% alcohol.
 - Covering your cough to prevent the spread of germs.
 - Asking family or friends not to visit when they don't feel well.
 - Staying in your room when sick.
 - Protecting yourself by getting vaccines for flu and pneumonia and encouraging others around you to do the same.
- Talk to your healthcare professional about steps you can take to stay healthy and prevent infections.

From: <https://www.cdc.gov/antibiotic-use/pdf/AU-nursing-home-trifold-brochure-P.pdf>. Accessed 9/1/24

Adopting Publically Available Educational Materials to Facility Need

Can Owners or Adults Responsible for Nursing Homes Top 10 Infection Prevention Questions to Ask a Nursing Home's Leaders

- 1. What infections commonly occur among residents in this facility?**
Why is this question important? Nursing home residents can have a higher chance of getting an infection due to facility items getting dirty, including items like handrails, bed rails, and other items that touch the body. A nursing home should be taking the common infections in its residents so they can be sure that actions are taken to reduce the spread of germs to residents.
- 2. When was the last outbreak (e.g., infection spreading among residents) in this facility?**
What was the cause (e.g., flu, norovirus, etc.)?
Did the facility request assistance from local public health authorities or any other partners?
Why are these questions important? Should having an outbreak or cases occur it is easier for a caregiver (like flu or diarrhea) from someone in great health through a nursing home. While some facilities require outbreaks, such as diarrhea and quickly starting, public health authorities can help limit the spread of the infection to new residents, staff and visitors.
- 3. How does the facility communicate with residents, family and visitors when an outbreak occurs?**
Why is this question important? Outbreaks of infections do occur in nursing homes. It is important the facility staff have a process for notifying residents, family members and visitors to ensure they take steps to decrease the chance of spreading the infection or getting it. Sometimes a facility might have to restrict visitors or stop new admissions temporarily in order to stop an outbreak. In these situations, it is very important that everyone is informed about what is happening.
- 4. Is the flu vaccine mandatory for all staff working in this nursing home?**
What percentage of your staff received the flu vaccine this year (or last year)?
Why are these questions important? A growing number of healthcare facilities are making it a requirement for staff as a measure to prevent patients and staff from flu. In nursing homes, getting a flu vaccine for staff can help reduce the spread of flu to residents. Even if the vaccine isn't mandatory for staff, all staff should be offered the vaccine and a nursing home should know how many of their staff received the vaccine during the last flu season. The nursing home should also know what percentage of residents received the flu vaccine during the last flu season.

From: <https://www.cdc.gov/longtermcare/prevention/antibiotic-stewardship.html>. Accessed 9/19/17.

Immanuel Fontenelle
an Immanuel care community

Top 10 Infection Prevention Questions to Ask a Nursing Home's Leader

- 1. What infections commonly occur among residents in this facility?**
As with most communal living typical infections include respiratory, gastrointestinal, and urinary.
- 2. When was the last outbreak in this facility?**
Last winter we had several cases of influenza.
- 3. How does your facility communicate with residents, families, and visitors when an outbreak occurs?**
3-1 communication with residents.
- 4. Is the flu vaccine mandatory for all staff working in this nursing home?**
Yes. The flu vaccine is mandatory for all staff unless they are allergic. If allergic they then have to wear a mask all flu season. Staff had a vaccination rate of 92.4% this last flu season. Residents had a vaccination rate of 92.9%.
- 5. If a staff member is sick, is he/she allowed to stay at home (or go home from work) without being on or time off?**
We encourage staff to stay home if not feeling well with signs and symptoms of any communicable disease. If a staff member shows up and is not feeling well they are asked to go home until they feel better.
- 6. How are facility staff trained to respond to questions about hand hygiene from residents and family?**
Education is provided to staff multiple times a year including computer based and skills check offs, along with verbal reminders of hand hygiene for compliance of practice.
- 7. Are residents with new diarrhea given separate toilet facilities until the cause of diarrhea is determined and/or the diarrhea is resolved?**
When a resident has diarrhea with unknown etiology, they are placed into transmission based precautions in their own room or a private room (if available) to protect them and the other residents until signs and symptoms have ceased for 72 hours or the physician has ordered anything different.

Courtesy of Chase Griffin, RN, Immanuel Fontenelle (now , Omaha, NE).

Antimicrobial Stewardship Education

nebraska icap asap

Nebraska ICAP & ASAP
@nebraska.icap.asap 2,234 subscribers 93 videos
The Nebraska Infection Control Assessment and Protection (ICAP) program...
icap.nebraskahealthcare.com and 1 more link

HOME VIDEOS PLAYLISTS COMMUNITY CHANNELS ABOUT

A Pandemic Within a Pandemic: COVID-19 Impact on Antibiotics
122 views · 1 month ago

2023 Nebraska Antimicrobial Stewardship Summit
122 views · 1 month ago

2023 Nebraska Antimicrobial Stewardship Summit ▶ Play all

Communication Strategies in Antibiotic Stewardship
Nebraska ICAP & ASAP
60 views · 3 months ago

Nebraska Healthcare Associated Infections and...
Nebraska ICAP & ASAP
75 views · 1 month ago

A Pandemic Within a Pandemic: COVID-19 Impact...
Nebraska ICAP & ASAP
122 views · 1 month ago

The Role of the MHE in Response to the Increase in...
Nebraska ICAP & ASAP
19 views · 1 month ago

The Gut Microbiome and Non-Traditional Therapies...
Nebraska ICAP & ASAP
39 views · 1 month ago

Fossilite Herding and Infectious Prevention...
Nebraska ICAP & ASAP
18 views · 1 month ago

2023 Nebraska Antimicrobial Stewardship Summit, Acute Care and Ambulatory Track ▶ Play all

https://www.youtube.com/@nebraska_icap_asap/featured

43

Summary

- ▶ Tracking, reporting, and education are important core elements that have direct impact on antimicrobial use
- ▶ Tracking includes policy compliance, antibiotic use, antibiotic resistance, and infection rates
- ▶ Reporting should inform prescribers, staff, residents of ASP activities and outcomes
- ▶ Education should be target audience specific and include goals of ASP, appropriate antibiotic prescribing/use

44

Free Online Resources

- ▶ Nebraska ASAP (asap.nebraskamed.com)
- ▶ CDC (cdc.gov/longtermcare/prevention/antibiotic-stewardship.html)
- ▶ AHRQ (ahrq.gov/nhguide/index.html)
- ▶ University of Rochester (rochesterpatientsafety.com/)