

OB HEMORRHAGE WORKSHEET

Medication	Route/Dosage	Considerations/Side effects	Time administered:
Oxytocin (1st line medication)	IV Run at 999 for 250 or 500mL (ask provider) IM 10mg	Usually no side effects n/v possibly	
Cytotec (misoprostol)	Rectal 800-1000 mcg (ask provider) Can be given PO/SL		
Methergine	IM 0.2 mg = 1mL Oral 0.2 mg for 24 hours (Q6-8HR)	Avoid hypertensive patients	
Hemabate	IM 250 mcg = 1mL	Avoid asthma patients Causes diarrhea → consider imodium prophylactically	
TXA (tranexamic acid)	IV 1000 mg = 10 mL over 10 mins	Push over 10 minutes to avoid hypotension	

Jada System (OB Intrauterine Suction Device Orderset!)	Bakri Balloon (OB Intrauterine Balloon Tamponade Orderset!)
<p>Remove air from seal prior to placement</p> <p>Suction to 80 mmHg</p> <p>Fill with 60-120 mL sterile water</p> <p>**Clamp tubing during transportation**</p> <p>If suction tubing full of blood = 100mL</p>	<p>Fill with sterile water per provider request</p> <p>Requires drainage bag</p> <p>Give antibiotics per provider request</p>

Dry weights:

Delivery Table Lab Sponge = 15 gms
 Sterile Blue Towel = 50 gms
 1 pink peri pad = 25 gms
 Small Red Biohazard Bag = 20 gms
 Big Blue COVIDIEN Chux = 105 gms

Small Blue Chux = 60 gms
 Perineal Cold Pack = 150 gms
 Ice Diaper = weigh before placing on patient for dry weight amount
 10 pack 4x4 Raytec Sponges = 20 gms